

Clark University

## Clark Digital Commons

---

Geography

Faculty Works by Department and/or School

---

3-15-2017

### Changes of forest cover and disturbance regimes in the mountain forests of the Alps

P. Bebi

*WSL - Institut für Schnee- und Lawinenforschung SLF - Davos*

R. Seidl

*Universität für Bodenkultur Wien*

R. Motta

*Università degli Studi di Torino*

M. Fuhr

*Université Grenoble Alpes*

D. Firm

*Univerza v Ljubljani*

*See next page for additional authors*

Follow this and additional works at: [https://commons.clarku.edu/faculty\\_geography](https://commons.clarku.edu/faculty_geography)



Part of the [Forest Sciences Commons](#)

---

#### Repository Citation

Bebi, P.; Seidl, R.; Motta, R.; Fuhr, M.; Firm, D.; Krumm, F.; Conedera, M.; Ginzler, C.; Wohlgemuth, T.; and Kulakowski, Dominik, "Changes of forest cover and disturbance regimes in the mountain forests of the Alps" (2017). *Geography*. 279.

[https://commons.clarku.edu/faculty\\_geography/279](https://commons.clarku.edu/faculty_geography/279)

This Article is brought to you for free and open access by the Faculty Works by Department and/or School at Clark Digital Commons. It has been accepted for inclusion in Geography by an authorized administrator of Clark Digital Commons. For more information, please contact [larobinson@clarku.edu](mailto:larobinson@clarku.edu), [cstebbins@clarku.edu](mailto:cstebbins@clarku.edu).

---

**Authors**

P. Bebi, R. Seidl, R. Motta, M. Fuhr, D. Firm, F. Krumm, M. Conedera, C. Ginzler, T. Wohlgemuth, and Dominik Kulakowski

Published in final edited form as:

*For Ecol Manage*. 2017 March 15; 388: 43–56. doi:10.1016/j.foreco.2016.10.028.

## Changes of forest cover and disturbance regimes in the mountain forests of the Alps★

P. Bebi<sup>a,\*</sup>, R. Seidl<sup>b</sup>, R. Motta<sup>c</sup>, M. Fuhr<sup>d</sup>, D. Firm<sup>e</sup>, F. Krumm<sup>f</sup>, M. Conedera<sup>g</sup>, C. Ginzler<sup>g</sup>, T. Wohlgemuth<sup>g</sup>, and D. Kulakowski<sup>a,h</sup>

<sup>a</sup>WSL-Institute for Snow and Avalanche Research SLF, Davos, Switzerland <sup>b</sup>University of Natural Resources and Life Sciences (BOKU), Vienna, Austria <sup>c</sup>Università degli Studi di Torino, DISAFA, Largo Braccini 2, 10095 Grugliasco (TO), Italy <sup>d</sup>Univ. Grenoble Alpes, Irstea, F-38402 St-Martin-d'Hères, France <sup>e</sup>Department of Forestry and Renewable Forest Resources, Biotechnical Faculty, University of Ljubljana, Slovenia <sup>f</sup>European Forest Institute, Freiburg, Germany <sup>g</sup>Swiss Federal Institute for Forest, Snow and Landscape Research WSL, Switzerland <sup>h</sup>Graduate School of Geography, Clark University, Worcester, MA, USA

### Abstract

Natural disturbances, such as avalanches, snow breakage, insect outbreaks, windthrow or fires shape mountain forests globally. However, in many regions over the past centuries human activities have strongly influenced forest dynamics, especially following natural disturbances, thus limiting our understanding of natural ecological processes, particularly in densely-settled regions. In this contribution we briefly review the current understanding of changes in forest cover, forest structure, and disturbance regimes in the mountain forests across the European Alps over the past millennia. We also quantify changes in forest cover across the entire Alps based on inventory data over the past century. Finally, using the Swiss Alps as an example, we analyze in-depth changes in forest cover and forest structure and their effect on patterns of fire and wind disturbances, based on digital historic maps from 1880, modern forest cover maps, inventory data on current forest structure, topographical data, and spatially explicit data on disturbances. This multifaceted approach presents a long-term and detailed picture of the dynamics of mountain forest ecosystems in the Alps. During pre-industrial times, natural disturbances were reduced by fire suppression and land-use, which included extraction of large amounts of biomass that decreased total forest cover. More recently, forest cover has increased again across the entire Alps (on average +4% per decade over the past 25–115 years). Live tree volume (+10% per decade) and dead tree volume (mean +59% per decade) have increased over the last 15–40 years in all regions for which data were available. In the Swiss Alps secondary forests that established after 1880 constitute approximately 43% of the forest cover. Compared to forests established previously, post-1880 forests are situated primarily on steep slopes (>30°), have lower biomass, a more aggregated forest structure (primarily stem-exclusion stage), and have been more strongly affected by fires, but less affected by wind disturbance in the 20th century. More broadly, an increase in growing stock and

\*Corresponding author. bebi@slf.ch (P. Bebi).

★This article is part of the special issue: "Ecology of Mountain Forest Ecosystems in Europe" published in the journal *Forest Ecology and Management* 388, 2017.

expanding forest areas since the mid-19th century have - along with climatic changes - contributed to an increasing frequency and size of disturbances in the Alps. Although many areas remain intensively managed, the extent, structure, and dynamics of the forests of the Alps reflect natural drivers more strongly today than at any time in the past millennium.

## Keywords

Land-use history; Secondary succession; Disturbance interactions; European Alps; Snow avalanches; Windthrow; Forest fire

---

## 1 Introduction

Mountain forests globally are undergoing major changes driven by factors related to climate, land-use, and natural disturbances (Dale et al., 2001; Kulakowski et al., 2012). While the understanding of all three of these driving forces has greatly increased, interactions between them are still difficult to disentangle. Natural disturbances such as fire, wind, insect outbreaks and avalanches are strongly affected by forest cover, forest structure, climate, and land-use (Kulakowski and Veblen, 2002; Seidl et al., 2011a; Kulakowski et al., 2011; Puerta-Piñero et al., 2012; Liu et al., 2015; Flatley et al., 2013). Especially in mountainous areas, which form a complex biophysical template for these disturbances and their drivers, the spatiotemporal complexities of associated dynamics are not yet well understood.

Forest cover has substantially increased since the 19th century in several mountain ranges of the world (Wear and Bolstad, 1998; Bunce, 1991; Kozak, 2003), mainly as a result of reduced or abandoned agricultural areas (Baldock et al., 1996; Gellrich et al., 2007). As past land-use can have multiple and long-term impacts on forest soils and successional pattern (Foster, 1992; Körner et al., 1997; Dambrine et al., 2006; Spohn et al., 2016), we can expect important differences between post-agricultural forests compared to areas that have long been forested (Flinn and Vellend, 2005; Foster et al., 2003). In addition to forest expansion, changes in climate and forest management have contributed to widespread changes in forest structure and biomass stocks. Across large parts of Europe, these changes have contributed to increased disturbances by wind, bark beetles, and wildfires over the past decades (Seidl et al., 2014; Wastl et al., 2013). However, in the specific context of the Alps it is not clear whether recent trends in disturbance regimes are primarily related to successional dynamics in newly established secondary forest, or to increasing biomass levels in previously established forests, and how strongly these changes are mediated by ongoing climatic changes. It is thus helpful to examine long-term trends to obtain better insights into the drivers of mountain forest dynamics.

Reliable data on forest cover changes and disturbance history is much more limited before the 19th century than for recent periods in the Alps, but several lines of evidence can provide insights into forest development and species shifts over longer time periods (Kaplan et al., 2009). In particular paleoecological data provide evidence of how forest composition and extent has changed during periods of increasing pressure of human land-use, and how fire regimes have changed in response to climate and land-use (Tinner and Kaltenrieder, 2005; Conedera et al., 2017). Evidence of past forest development and the historic variation of

different disturbance regimes is often provided by dendroecological reconstructions of disturbance regimes (e.g. Janda et al., 2017; Panayotov et al., 2017), but in some regions the influence of human management over past centuries was strong, obscuring the evidence of natural disturbances (Kulakowski and Bebi, 2004). Empirical data can be complemented with simulation models that can help to elucidate current and future patterns and composition of mountain forests, and how they are affected by climate, land-use and natural disturbances (e.g. Temperli et al., 2013; Thom et al., 2016). In contrast to the forest history since the 19th century, which has been characterized by increasing biomass and disturbances (cf. Usbeck et al., 2010a, 2010b, Appendix, Table A1), we have a relatively fragmentary picture of the processes that contributed to the massive decreases of forest cover and biomass prior to the 19th century (Kaplan et al., 2009; Küster, 2010). Consequently, our long-term understanding of the variability in disturbance regimes remains cursory for forest ecosystems such as those in the European Alps, which have a long and intensive management history (Bätzing, 2003; Mathieu et al., 2016). As a result, no long-term and broad-scale overview on natural disturbance regimes of the Alps exists to date.

In this contribution we briefly synthesize the available information on long-term (i.e. >100 years) forest cover changes and disturbance regimes in the Alps. We combine this information with a compilation of forest inventory and forest disturbance data for the entire mountain range of the Alps, as well as detailed data on forest cover, structure, and disturbance development since the 19th century for the Swiss Alps (Ginzler et al., 2011). Based on these sources of information we put recent forest structure and dynamics into a long-term context and we address the following main questions: (1) What are the recent trends in forest cover, structure, and disturbance regimes since the 19th century, and how do they relate to the long-term context of forest development? (2) How do secondary forests that established since the 19th century differ from pre-existing forests in terms of stand structure and natural disturbance regimes?

## 2 Long term forest composition and land use changes

The European Alps extend over approximately 1000 km, from the French and Italian Mediterranean coast across Switzerland, southern Bavaria, northern Italy, Austria and Slovenia, and have a total population of 14 million people (Chartré et al., 2010). The mountain peaks reach elevations of >4000 m a.s.l. and are intersected with deep valleys, some of which are >100 km long and divide the mountain range into major massifs. The Alps are a relatively young mountain system, whose “step-like” morphology was contoured by the Pleistocene glaciation. Bedrocks can be divided into calcareous and crystalline material. The climate is characterized by strong environmental gradients ranging from oceanic to dry climate. The most widespread forest types are mixed European beech (*Fagus sylvatica* L.) and silver fir (*Abies alba* Mill.), pure Norway spruce (*Picea abies* (L.) H. Karst.), and mountain pine (*Pinus mugo* Turra s.l.) in the front ranges, while European larch (*Larix decidua* Mill.), Swiss stone pine (*Pinus cembra* Mill.) and Scots pine (*Pinus sylvestris* L.) may replace them in the dry central Alps (see Conedera et al., 2017 for more details).

Early changes in forest cover and forest composition since the late glacial-Holocene transition have been reconstructed based on paleoecological records (Kral, 1995; Tinner and

Kaltenrieder, 2005; Conedera et al., 2017). These records show evidence of a relatively rapid invasion of pioneer species like European larch, and different pine species, occurring as early as 11,400 years before present (y BP) and extending as far as the current subalpine belt (Blarquez et al., 2009; Tinner and Kaltenrieder, 2005). Major current tree species of the Alps like silver fir (glacial refugia in the south), Norway spruce (refugia in the east), and European beech (different refugia in the south, west and east) immigrated to the Alps after 9000 y BP (Kral, 1995; van der Knaap et al., 2005).

Land-use has influenced forest dynamics at least in parts of the Alps since ca. 7500 y BP, when Neolithic herdsman started using fire to expand pastures for grazing in mountain forests (Conedera et al., 2017; Winckler, 2012). These early human impacts and the intense use of fire have not only changed forest cover and forest density in some regions of the Alps, but have also shifted species composition. For example, anthropogenic fires combined with successive intensive browsing facilitated expansion of *P. abies* into areas formerly occupied by *A. alba*, which is more sensitive to such disturbances (Schwoerer et al., 2015; Conedera et al., 2017). A second (between 5000 and 3500 y BP) and a third (between 1200 and 700 y BP) wave of increase in the human population and human migration into the Alps led to permanent settlements in higher elevation areas, resulting in major and wide spread human impacts on mountain forests (Schuler, 1988; Winckler, 2012). Accelerated slash and burn management during this third wave reduced forest cover in central Europe, which was already below or comparable to current levels (Hauser, 1964; Bork et al., 2001).

With an increasing awareness of trade-offs between deforestation and the occurrence of natural hazards, the first written regulations against further exploitation of protection forests (i.e., forests that protect against natural hazards) were enacted in the 13th and 14th centuries (Price, 1988). In spite of such regulations, people continued to intensively exploit mountain forests and their products for energy and construction materials as well as extracting litter and pasturing over the following centuries (Mathieu et al., 2016). Deforestation and exploitation were partly slowed due to the Black Death and the ensuing decline in the human population in the 16th and 17th century (McEvedy and Jones, 1978). There is strong evidence of accelerated deforestation across most of the Alps for the late 18th and early 19th century, when human population again increased and the demand of wood resources was strongly boosted by industrialization (Mather et al., 1999; Bätzing, 2003).

Multiple factors acting simultaneously finally halted forest exploitation in the Alps in the mid-19th century, with the importance of these factors varying from region to region. In some areas, such as the French Alps and the SW parts of the Italian Alps, regional depopulation (partly amplified by unfavorably cool climatic conditions) resulted in an expansion of forest cover (Bätzing, 2003; Motta et al., 2006b). In contrast, heavy flooding (especially in the 1860s) and a generally increasing awareness of the protection effect of mountain forests against natural hazards contributed to afforestation, stricter laws, and adapted management in mountain forests of the Northern Swiss and Bavarian Alps (Mather and Fairbairn, 2000), the Austrian and Slovenian Alps, and parts of the French Alps (Sonnier, 1991). Additionally, the combination of increasing agricultural efficiency and a gradual replacement of fuel-wood by coal and other fossil energy sources decreased the pressure on forests towards the end of the 19th century (Mather and Fairbairn, 2000). The

increasing globalization (e.g. transport, trade) also contributed to a reduced use and exploitation of forests, which in turn contributed to denser forests and larger areas with progressing secondary succession (Fig. 1).

### 3 Long-term forest disturbance regimes

Natural disturbances (i.e. discrete large pulses of tree mortality from agents such as avalanches, fires, winds or insect outbreaks) have been important drivers of the ecology of the mountain forests of the Alps for millennia. Snow avalanches and other snow-related disturbances have been integral in the mountain forests of the Alps, which is evidenced by (1) reconstructions of the snowline during the Holocene (Patzelt and Bortenschlager, 1973), (2) long-term historical avalanche records of the past centuries (Latenser and Pfister, 1997), and (3) adaptations of tree species in snow-rich environments to snow related disturbances (Bebi et al., 2009). As steep mountain peaks in large parts of the Alps exceed the current elevation of the treeline (1750–2350 m a.s.l., Paulsen and Körner, 2009) and accumulate large snow packs during winter, snow avalanches frequently intersect forested areas in the Alps and contribute to a heterogeneous forest structure (Bebi et al., 2009; Vacchiano et al., 2015). Extensive areas that had been deforested since the beginning of human settlement in the Alps increased avalanche frequency by creating new release areas in formerly forested terrain resulting in a long-term absence of forest cover due to repeated avalanche disturbance (Küttel, 1990).

In addition to avalanches, snow breakage across different spatial scales has been a widespread disturbance in the forests of the Alps (Coaz, 1887; Weigl, 1997; Rottmann, 1985; Klopčič et al., 2009, Appendix, Table A1). Damage to trees that are not well adapted to large snow loads can be considerable, particularly in mid-elevation areas and/or during exceptionally wet and heavy snow events (Hlasny et al., 2011a; Deankovic, 1969). Although different attributes of stand structure may modulate the susceptibility to snow breakage, the long-term variation in disturbance from snow breakage is, like for avalanches, mainly driven by climate.

Fire regimes in the Alps are highly heterogeneous, with a clear difference between relatively large burned area and high fire frequency in the Southern Alps, due to a convergence of environmental and climatic factors that enhance fire ignition and spread, and lower fire frequency in the Northern Alps (Fig. 2, Appendix, Table A1, Wastl et al. (2013)). In contrast to fire activity on the southern slopes of the Alps, which peak in the relatively dry winter season and which are mainly influenced by human ignitions, forest fires are most frequent in summer in Central Alpine valleys, which are characterized by persistent snow cover, dominance of coniferous forests (especially highly ignitable Scots pines, *Pinus sylvestris* L.), and relatively high proportion of lightning-caused forest fires (Cesti et al., 2005; Conedera et al., 2006). An elevated probability for lightning-caused fires also exists along the eastern and south-eastern rim of the Alps in Austria (Müller et al., 2012). Besides such environmental and climatological influences and their changes over time, human activity has been, by far, the main cause of fires for ca. 7500 years in many regions of the Alps (Conedera et al., 2017; Arndt et al., 2013). Slash and burn management decreased forest cover in many parts of the Alps and shifted tree species composition. This effect is

particularly evident in the Southern Alps, where human-caused fires strongly reduced the abundance of fire-sensitive species (Tinner et al., 1999). Management by fire also played an important role in the Northern Alps during the same time period, but fire frequencies were too low to result in the disappearance of entire forest communities (Tinner et al., 2005). Where natural fire frequency was historically high (e.g. in dry zones of the Central Alps), forest vegetation has been more fire adapted and changed less in response to human-induced increases in fire frequencies (Stahli et al., 2006). Since Roman Times (ca. 2000 y BP), fire use has been less extensive, but has persisted as an important ecosystem management tool and as a driver of landscape change in many regions of the Alps (Conedera et al., 2017; Valese et al., 2014).

Data about disturbance from windthrow and bark beetle outbreaks stem mostly from records going back only to the mid- 19th century (Usbeck et al., 2010a, 2010b; Bottero et al., 2013; Seidl et al., 2014, Appendix, Table A1), and suggest strong spatiotemporal variations in occurrence and extent. In the Alps strong winds with potentially damaging effects on forests occur mainly due to westerly or northerly winter storms (Gardiner et al., 2010; Usbeck, 2015). Consequently, exposed forests of the Northern and North-Western Alps are more prone to large-scale wind disturbance compared to mountain forests in the inner Alpine valleys and the Southern Alps, which are generally protected from strong winds by the northern front range (Usbeck et al., 2010a, 2010b). However, foehn winds can also result in large wind disturbances in the central and eastern parts of the mountain range. Bark beetle outbreaks are highly correlated with spruce abundance and standing volume, growing season temperature, drought stress, and preceding disturbances of which windthrow is the most important, as it creates large amounts of virtually defenseless breeding material for beetle development (Thom et al., 2013; Stadelmann et al., 2014). Historical records on windthrow events and bark beetle outbreaks before the 19th century are widely missing (Pfister, 1988) and dendroecological reconstructions of these disturbance events are difficult in forests with intensive management history (Kulakowski and Bebi, 2004). Our knowledge of the historical range of variability of these disturbance agents in the Alps is thus very limited compared to other European mountain ranges (Vacchiano et al., 2016; Kulakowski et al., 2017), and can only vaguely be deduced from a relative short observation period and from analogies with similar mountain forest ecosystems.

#### 4 Methods of analyzing forest cover dynamics since the 19th century

To analyze forest dynamics in the Alps, we compiled and analyzed available records on forest area, forest structural change (including living and dead timber volume, as well as the growing stock of relevant tree species) and available information on occurrence, size and damage pattern of major natural disturbances for all alpine countries (Austria, Germany, France, Italy, Liechtenstein, Slovenia, Switzerland). Changes of forest area and other attributes were computed as average rates of changes per decade based on the back-calculated values at the beginning of each decade.

In addition, we compared forests that established after 1880 with pre-existing forests and analyzed forest cover changes since 1880 under different topographical settings in the Swiss Alps, where these data were available. To do so, we intersected digitized historical forest

cover maps for the years 1880, 1940 and 2000 (Ginzler et al., 2011) with plots of the Swiss National Forest Inventory (NFI; regular 1.4 km grid), excluded all plots that have not been forested according to the NFI-data of 2004–2006 or the forest cover map of the year 2000, and derived forest presence/absence between 1880 and 2000 separately for the Northern Prealps, Central Alps and Southern Alps. NFI stand structural data were recorded (1) by field crews in two concentric circles with sizes of 200 m<sup>2</sup> (inner circle) and 500 m<sup>2</sup> (outer circle), and (2) for an area of 50 \* 50 m surrounding the study plot, where additional variables on stand structure and management history were assessed based on aerial photo interpretation, field surveys and surveys by regional foresters (Keller, 2005).

We used GIS (*ArcGIS* version 10.1) to define the distance from the potential treeline as well as the presence of wind and fire disturbances in these plots. The potential treeline was calculated following the approach of Paulsen and Körner (2009), where the regional treeline was deduced from highest forest patches identified by a moving window algorithm. In order to compare the occurrence of windthrow and fire in secondary (post-1880) versus pre-existing (pre-1880) forests we applied a spatial overlay analysis of the historic maps with digitized disturbance data. We derived wind disturbances from remotely sensed polygons of forest damage data from the winter storms Vivian (in 1990) and Lothar (in 1999) (Usbeck et al., 2012) and intersected these polygons with NFI-plots and the appended information on current and 1880-forest cover for each NFI-plot. The digitized inventory of 487 fire ignition points (fire plots) of the Canton of Grisons between 1971 and 2015 (Pezzatti et al., 2010) was directly intersected with digitized historical maps of 1880. For the intersected points we calculated then the association between the expected disturbance probability (number of disturbance plots under uniform distribution over the two forest categories) and the observed number fire- and windthrow plots in pre-1880 versus post-1880-forests based on Chi-square statistics. The intersections between NFI-plots and the 1880-map were also used to test for associations between the two forest categories and a number of other known attributes for each NFI-plot including variables on current forest structure and past forest management (Abegg et al., 2014). Pearson's Chi-squared tests (with Yates' continuity correction) were used to test associations with binary variables and Wilcoxon-Mann-Whitney tests for associations with continuous variables (R core Team, 2013).

## 5 Results on forest cover dynamics since the 19th century

### 5.1 Forest trends and disturbance regimes in the Alps

All available data sets – both ground inventory and repeated forest cover maps based on remote sensing – across all regions of the Alps indicate an increase in forest cover in recent decades. However the rates of forest area change since the availability of reliable data in the late 19th or early 20th century varied considerably, both spatially and temporally, and range from +7 to +150% overall during 23–176 years (Fig. 3). Forest area expanded most rapidly in parts of the Italian Alps (e.g. +8.7% per decade since 1962 in Val Masino, Sondrio (Martelletti et al., unpublished data), +6.2% per decade since 1954 in Val Vigezzo, Verbania (Vacchiano et al., 2008)), in the Southern Swiss Alps (average rate of +7.3% per decade since 1880), and in the Austrian province of Salzburg (+7.0% per decade since 1928, BFW, 2016; Weigl, 1997). Forest cover expanded less rapidly in Bavaria (Germany; +0.7% per

decade since 1900), in the French Prealps (+3.5% per decade between 1850 and 1990) and the Northern Swiss Alps (+2.1% per decade between 1880 and 2013). On average (mean of all reported values), forest area across the Alps increased by +3.7% per decade since 1930 and +4.3% per decade since 1990.

Live and dead biomass also strongly increased since the end of the 19th century and continues to increase. Based on inventory data of different countries in the Alps (since 1973 for the Northern French Alps, since 1992/96 for Austria and since 1993/95 for the Swiss Alps), the total live wood volume increased in all regions of the Alps, averaging +9.7% of growing stock per decade, and ranging from +0.4% per decade in the northern Swiss Alps to +16.9% per decade in the eastern Slovenian Alps and to +17.1% per decade in the Southern Swiss Alps (Table 1). Dead wood volume increased even more in the same time period, with rises between +39.3% and +105.8% per decade (Table 1). Shifts in species composition differ across the Alps, and include an increased share of silver fir on the total growing stock in parts of the French Alps (+11.0 % per decade) that contrasts with a decrease in parts of the Eastern Alps (e.g. -2.6% per decade in Vorarlberg (Austria) and the eastern Slovenian Alps and -3.8% in Liechtenstein, Table 1).

The occurrence, size and damage volume of recorded natural disturbances varied strongly by disturbance agents and regions within the Alps (Appendix, Table A1). Overall, the greatest reduction of forest growing stock was associated with large windthrow events, which damaged an average of >2.5 millions of cubic meters per year since 1990 in the Northern Alps and resulted in small (scattered canopy openings) to large (>50 ha) patches of windthrow, which was often followed by salvage harvesting. Snow and ice breakage events also varied from very small (<0.1 ha) to substantial (1.5 mio m<sup>3</sup> damaged in the extreme ice breakage event in the Slovenian Alps in 2014), but could not be distinguished clearly from other abiotic factors as wind events in all areas. Bark beetle disturbance has been quantified since the early 20th century and the recorded timber volume damaged per year has strongly increased (up to >2 mio m<sup>3</sup>/yr) in recent decades. In comparison with other disturbances, avalanches often resulted in relatively large disturbed patches (maximum of 93 ha and a median of 1 ha), but the damaged timber volume (<0.2 mio m<sup>3</sup>/yr in the Swiss Alps even in extreme avalanche winters) was comparatively low. Fire disturbances, which increased in frequency from the Northern Alps to the Central and the Southern Alps, burned over several 100 ha in extreme events, but more than half of all fires were smaller than 0.3 ha in size (Appendix, Table A1).

## 5.2 Detailed analysis of forest change in the Swiss Alps

Forest area change varied across natural gradients: increases were greatest on slopes >30° compared to more gentle slopes, in particular in the Central Alps, where >90% of the forest area increase occurred on slopes with a steepness of >30°. The highest rates of increase since 1880 occurred near the treeline ecotone (0–200 m below the potential treeline), but also areas that are >800 m below the current potential treeline changed considerably (Fig. 4b).

Secondary forests (i.e. areas not identified as forests in the 1880 maps) make up ca. 43% of all current forests in the Swiss Alps, and differed distinctly from pre-existing (pre-1880)-

forests in the most recent forest inventory (Table 2). Post 1880-forests have lower biomass (both living and dead stock volume), are more often characterized by low dominant diameters (between 10 and 30 cm DBH) and less represented in large dominant diameter (>50 cm DBH) classes. In spite of still lower biomass, the canopy closure in secondary forests was more often classified as closed (dense forest with competing crowns) or as spatially aggregated (interspersed dense patches within the 50 × 50 m plots), but less often as vertically heterogeneous. Secondary forests more often showed signs of grazing, but were otherwise less often actively managed over the last 50 years. Spruce, fir, and beech were more often the dominant tree species in pre-existing forests, while larch and miscellaneous broadleaved species (e.g. *Acer pseudoplatanus* L., *Sorbus aucuparia* L., *Alnus viridis* (Chaix) DC) were more likely to dominate in secondary forests.

Forests that established after 1880 were less affected by the two largest storms of the 20th century, namely Vivian (in the year 1990) and Lothar (in the year 1999), compared to pre-existing forests ( $p = 0.005$ , Chi-square test, Fig. 5). Based on a record of 487 fires in the Swiss Canton Graubünden the density of fires that occurred between 1971 and 2015 was higher in secondary forests ( $p = 0.00013$ , Chi-square test) (Fig. 6).

## 6 Discussion

Our review of forest changes in the Alps prior to the 19th century and our analysis of spatial data since the 19th century indicate that changes in forest area and forest composition during the last millennium have been strongly related to changes in land-use. Forest area across the Alps has dramatically increased during the last century at approximately + 4% per decade after a minimum during the mid-19th century. The rates of forest area change have varied considerably in response to regional socio-economic and environmental factors (Bätzing, 2003; Gellrich et al., 2007), but a general increase is evident across the entire European Alps. While both live and dead biomass have increased across different regions of the Alps (Table 1, Abegg et al., 2014, BFW, 2016), changes in species composition follow more regional patterns and are influenced by a variety of factors. For example, regionally diverging trends in silver fir share have been attributed to different levels of ungulate browsing (Ammer, 1996; Didion et al., 2009; Klopčič and Boncina, 2011), while strong expansion of the same species in the French Alps (Chauchard et al., 2010) and of European larch in the western Italian Alps (Motta et al., 2006b; Garbarino et al., 2011) have been mainly attributed to an extensification of live-stock grazing during the 20th century.

Furthermore, our in-depth analysis of forest area change in the Swiss Alps, comparing maps from 1880 to more recent maps (Ginzler et al., 2011), confirmed strong regional differences as well as the decisive role of topography and land-use history for forest development (Fig. 4). Major increases in forest area before 1940, particularly near the upper treeline, may in part be artifacts of slightly different criteria for the assessment of forest cover in the different maps analyzed here (Ginzler et al., 2011), but are likely to reflect decreasing land-use intensity, which began first at the most remote high elevation sites. This is consistent with other research showing that forest expansion near treeline in the Alps during the late 20th century has mainly been a bounce-back from a treeline depressed by previous anthropogenic activity, rather than a climatically induced advance of the treeline (Gehrig-Fasel et al., 2007).

This further suggests that in comparison to the dominant effects of land-use, warmer temperatures during the last decades probably had a relatively minor effect on the expansion of forest cover since the end of the 19th century (Kulakowski et al., 2011).

The specific land-use history of the Alps is also reflected in the current forest structure of secondary forests that established after 1880 compared to pre-existing forests. After the decrease in grazing pressure, these forests are either still in an early stage of stand initiation (apparent in the higher proportion of scattered and clustered forest structures) or they are already in more dense stages of forest development with increasing competition and stem-exclusion (Krumm et al., 2012). The initial conditions for these secondary forests were characterized by a high availability of light, exposed mineral soil in grazed areas, and an absence of structural heterogeneity typical for naturally disturbed sites (e.g., remnant large trees, logs, and pit-and-mound topography). Long-lasting legacies of past land-use also have been reported for soil nutrient concentrations (Spohn et al., 2016) and seed banks (Plue et al., 2008). The timing of forest establishment likely influenced also the species composition. Late successional species are more abundant in pre-1880 forests, while early successional and light-demanding species (*Larix decidua*) occur more frequently in post-1880 forests (Table 2). However, such differences between pre- and post-1880 forests have to be interpreted carefully because of differences in environmental drivers, and also because most of the pre-1880-forests of the Alps have a long-term management history.

In contrast to our increasingly clear view of the large-scale patterns of forest development after 1880, temporal development of forest cover changes before the 19th century can only be cursorily characterized based on available historical records and paleoecological data. The potentially large variation in the timing and intensity of anthropogenic forest area change and forest exploitation across the Alps prior to 1880 is difficult to show based on available evidence. Nevertheless, studies suggest that intensive land use has affected most forests in the European Alps, has strongly reduced forest area and created open forest structures by the late 18th century (Landolt, 1862; Kaplan et al., 2009). Similarly, decreasing biomass until the 19th century and a subsequent fast recovery of biomass also has been documented for forests outside of the Alps which were subject to similar changes in land use (e.g. Foster, 1992; Mather et al., 1999). The effects of forest cover changes prior to the 19th century on historical disturbance regimes are difficult to assess, not only because of uncertainties in timing and intensity of forest cover changes but also because of obvious difficulties in reconstructing natural disturbances in intensively managed forest landscapes (Kulakowski and Bebi, 2004). Our knowledge about the historical range of variability of disturbance regimes in the Alps (cf. Kulakowski et al., 2017) has thus to be deduced from a synthesis of (1) disturbances occurring since the 19th century (summarized in Table A1) and (2) our fragmentary picture of forest cover change and disturbance regimes before this forest transition.

## 6.1 Windthrow

Forests that established after 1880 have been less affected by recent windthrows in 1990 and 1999 compared to older forests. This may be explained by higher susceptibility of older, pre-existing forests with taller trees and higher biomass (Kulakowski and Veblen, 2002;

Gardiner et al., 2010; Thom et al., 2013) and suggests that the observed increase in storm damage during the 20th century across Europe (Schelhaas et al., 2003) could be explained more by the aging of pre-1880 forests and less so by the establishment of new forests (see also Seidl et al., 2011a). At the same time, however, storms with critical wind speed have become more frequent since the 1940s (Usbeck et al., 2010a), with particular consequences for forests with high stock rates (Usbeck et al., 2010b). Alternatively, given that post-1880 forests are located preferentially in topographic settings that were preferable for agriculture, lower damage levels in recent wind storms may also reflect differences in topographic exposure. Thus, future research should test whether differential wind damage in pre- and post-1880 forest is due to differences in forest structure or topographic exposure.

Because of lacking data, assessments on storm damage before the 19th century remain difficult. However, extratropical cyclones have likely led to winter storms that are similar to those that affect the Alps today for a long time (Kraus and Ebel, 2003), and historical wind damage in other mountain ranges of Europe (Kulakowski et al., 2017) also suggest that wind disturbances were likely common before significant human influence. Based on known relationships between growing stock, forest structure, and storm damage (Dobbertin, 2002; Gardiner et al., 2010) we can assume that historic land-use and management in the Alps have partly reduced the effects of wind disturbances, but that wind has been among the most relevant forest disturbance agents long before the 19th century.

Whether wind disturbance – currently the single most important disturbance agent in the Alps – will cause even more damage in the future remains highly uncertain, not least because projections of future wind dynamics remain challenging. However, with mountain forests responding to longer growing seasons, higher mean temperatures and, to some extent, CO<sub>2</sub> fertilization, increasing stocking levels can be expected to make forests more prone to wind disturbance (Seidl et al., 2011b). Furthermore, as most wind disturbances in the Alps occur in winter, and as tree stability is highly sensitive to soil frost (Usbeck et al., 2010a, 2010b), a decreasing period of soil frost could further increase forest susceptibility to wind in the future.

## 6.2 Avalanches and snow breakage

Forest expansion in the Alps since the 19th century has mostly occurred on steep slopes above 30°, and has thus led to a decreased avalanche activity in many areas (Bebi et al., 2009). This recent trend has partially compensated for the increasing avalanche activity due to deforestation and forest degradation before the 19th century. However, compensation of former increases in avalanche disturbance has not occurred where anthropogenic deforestation (partly combined with climatic shifts during the Little Ice Age) allowed the development of new avalanches, which continue to disturb exposed forests and inhibit their growth and development. For example, reforestation of large parts of the Urseren Valley (Switzerland), where the original forest had been reduced to four small dispersed fragments by a combination of land-use and avalanches, has not been possible under the current conditions of climate and avalanche disturbances (Föhn, 1978), even though paleoecological records indicate that this valley was forested during most of the Holocene (Küttel, 1990). Warmer winter temperatures and decreasing days with minimum snow depth required for

avalanche will probably further reduce the importance of avalanche disturbances in forested terrain in the future, and will further promote shifts from dry avalanche regimes to wet avalanche regimes (Castebrunet et al., 2014). However, where avalanche release zones are above the current treeline and forests are shaped preliminary by recurring avalanche disturbance, avalanches will likely continue to disturb forests - as they have throughout most of the Holocene.

Forest structural characteristics such as large proportions of pole stage stands and high h/d-ratios increase susceptibility to snow breakage (Rottmann, 1985; Nykänen et al., 1997; Hlasny et al., 2011a). Such structural characteristics are typical for stands that established after 1880 (Table 2) and are likely to become more widespread with increasing crown closure and competition in these stands (Krumm et al., 2012). However, snow breakage events are primarily related to specific weather events, and climate warming is likely to reduce the frequency of snow fall events in forested areas of the Alps (Schmucki et al., 2015), which will probably compensate for potentially increased susceptibility to snow breakage due to changes in forest structure.

### 6.3 Insect outbreaks

Based on current knowledge - which is mainly derived from data on the time since the 19th century - bark beetles (and here primarily *Ips typographus*) are the most relevant biotic disturbances in the mountain forests of the Alps. Beetle outbreaks often follow wind disturbances and have similarly increased over the last decades, mainly as a function of increasing volume of potential host trees (i.e. mainly *P. abies*) and increasing summer and winter temperatures (Seidl et al., 2014; Stadelmann et al., 2014). However, in contrast to evidence of regular outbreaks of e.g., larch budworm (*Zeiraphera diniana*), obtained from dendroecological reconstructions (Büntgen et al., 2009), we have almost no direct information about bark beetle outbreaks prior to the 19th century. A detection of earlier outbreaks with dendroecological methods (Cada et al., 2016) is not reliable in the Alps because of the overwhelming influence of forest management. Historical documents for periods prior to the 19th century are hardly specific enough to clearly identify bark beetle outbreaks (Pfister, 1988). Because of missing information and interactions between different drivers of beetle outbreaks assumptions about the historical range of variability of bark beetle disturbances remain challenging. However, based on information on the rigorous exploitation of living and dead biomass in mountain forests in earlier periods as well as analogies with similar mountain ecosystems in Europe (Kulakowski et al., 2017) we can assume that bark beetle outbreaks are part of the natural disturbance regime in spruce dominated forests in the Alps. Furthermore, it is likely that bark beetle populations (and consequently also the population of their predators) have been kept below their natural levels during periods of most intensive forest use before the 19th century.

In the face of increasing temperature, high and still increasing volume stock of *P. abies* at high elevations, and the aging of relatively young and even-aged forests stands, further increases of bark beetle activities in the future must be expected (Seidl et al., 2009). This may be particularly relevant at higher elevations where increasing temperatures will make new areas of spruce forests susceptible to bark beetles (Hlasny et al., 2011b). Furthermore,

an increasing propensity for drought events has the potential to trigger bark beetle outbreaks more frequently in the future (Netherer et al., 2015; Seidl et al., 2016). Warmer and drier conditions can also amplify the interaction between wind and bark beetle disturbance, further increasing future disturbances in the mountain forests of the Alps (Seidl and Rammer, in press). Over longer time scales, however, a climatically-induced decrease of Norway spruce at lower elevations may provide a negative feedback on bark beetle outbreaks (Temperli et al., 2013).

#### 6.4 Fires

The increase in forest area and biomass since the 19th century has resulted in higher continuity and extent of fuels. Retrospective analysis from the 20th century showed that an increase of fire frequencies in the Southern and Central Alps was at least partly related to an increase in fuel availability and fuel continuity due to a decrease of agricultural land-use in lower elevations (Zumbrunnen et al., 2009; Conedera et al., 2017), whereas warming and earlier spring seem to be more important in higher elevations and regions with less change in forest-cover (Zumbrunnen et al., 2009; Westerling et al., 2006).

We found higher fire densities in secondary forests that established after 1880 compared to pre-existing forests. This suggests that increased forest cover and connectivity between potentially burnable forest patches may have contributed to the observed increase of forest fires in the Alps (Zumbrunnen et al., 2009). These relationships vary strongly across different elevational zones and forest types, and may also be influenced by location of forests and human factors such as firefighting techniques or fire-inducing activities at the wildland-urban interface (Conedera et al., 2015). Thus, future research should test whether the increase in fires in these younger forests is due to the structure of the forests, or rather a factor of their location.

For the future we can expect that the ongoing forest cover increase and build-up of biomass (Table 2, Abegg et al., 2014) will continue to provide fuels for potential fires, which will be especially important where fires are limited by the abundance or continuity of fuels. Furthermore, while the complex topographical template influencing fire regimes in mountain regions will remain constant, increasing temperatures, summer droughts, and shorter snow duration will increase the probability for forest fires in the Alps (Arpaci et al., 2015), even in relatively mesic forests in which fires have not been historically important. Particularly in areas where forest vegetation is less adapted to frequent fires (Tinner et al., 2005), this may lead to drastic vegetation shifts (Moser et al., 2010). In the Southern Alps, we also can expect an increase in the fires at higher elevations and in forest types that until now have only been marginally affected by fires (Ascoli et al., 2013).

#### 6.5 Interactions among disturbances

Studying interactions among different disturbances is challenging, but of high ecological and societal relevance because disturbance interactions may lead to unexpected, rapid, and nonlinear changes in ecosystems (Paine et al., 1998; Buma, 2015), especially under a changing climate (Kulakowski et al., 2012, 2013). One of the most important disturbance interactions in the Alps involves positive feedbacks between the accumulation of fresh

deadwood and subsequent insect outbreaks (Bottero et al., 2013; Stadelmann et al., 2014). Other positive feedbacks can result from the removal of biomass due to windthrow, fire, or other disturbance and subsequent gravitational hazards like avalanches, rockfall, or shallow landslides. Such interactions are of particular societal importance in the densely populated Alps, and more knowledge is needed on interactions between a broad range of disturbance agents, in different forest types, and under different management regimes (Conedera et al., 2003; Bebi et al., 2015).

Negative feedbacks between disturbances, leading to lower susceptibility to subsequent disturbances, reduced biomass, and a more fragmented forest cover have received less attention in the Alps. The most obvious examples here are avalanche tracks, which may act as fire breaks (Veblen et al., 1994). Additionally, negative feedbacks between the reduction of biomass through windthrow, fire, insect outbreaks or snow breakage, and the susceptibility to subsequent disturbance (e.g. Kulakowski and Veblen, 2002; Kulakowski et al., 2003) may become increasingly important.

While disturbances which open the canopy generally reduce the dominance of the prevailing species and increase species diversity (Wohlgemuth et al., 2002; Vacchiano et al., 2016), this effect may be dampened by interactions with ungulate browsing, which may selectively exclude browsing-sensitive species from establishing and facilitate spruce dominance (Didion et al., 2009; Rozman et al., 2015; Conedera et al., 2017). In contrast, post-disturbance coarse woody debris may offer physical protection and thus reduce browsing pressure on tree regeneration (Bottero et al., 2013). In spite of such often neglected interactions between canopy removing and other disturbances, the overall effect of individual and interacting disturbances will probably counteract the general trend of growing biomass across the Alps. The resulting heterogeneity and lower biomass may reduce the risk of even larger disturbances in future. This mechanism may be particularly important in the light of climate change, as disturbances can present opportunities for ecological communities to adapt to new conditions by allowing new, better adapted species to establish (Buma and Wessman, 2013). The importance of both positive and negative feedbacks between different natural disturbances may thus considerably increase in future and should be emphasized more in the future management of mountain forests ecosystems in the Alps.

## 7 Synthesis and conclusion

Forest changes in the European Alps have been strongly driven by land-use, both before and after a major transition from forest loss to forest gain in the 19th century. The long-standing effects of land-use have had, and continue to have, a strong influence on forest dynamics and the disturbance regimes of mountain forests in the Alps. Human-induced decreases of forest area and density, peaking in the early 19th century, created new avalanche release areas in parts of the Alps. Furthermore, extraction of biomass and the resulting lower stocking levels reduced large-scale disturbances by fire, bark beetle and windthrow before the 20th century. The extensive forest areas in the Alps established after the 19th century on former agricultural land are currently characterized by relatively young stands with a high potential for further biomass accumulation and homogenization of forest structure. Susceptibility to disturbances by bark beetle, fire, and wind is likely to further increase in these areas over the

coming decades, particularly in combination with the climatic changes which are expected for the future.

The future management of mountain forest ecosystems in the Alps has to take into account the important and potentially increasing influence of natural disturbances under climate warming, while considering the particular and long-lasting effects of land-use history. It is not possible and, from an ecological perspective, also not desirable to impede these natural disturbances. However, where the protection of forests against natural hazards or other ecosystem services are threatened by disturbances and other natural processes, management may focus on reducing risks and increasing the resilience of mountain forests (Seidl, 2014). This can be achieved by disturbance management that allows forests to adapt to future environmental conditions and by counteracting the growing biomass and reduced fragmentation, particularly in secondary stands established after the mid-19th century.

Because of a strong and widespread anthropogenic effect on forest dynamics in the mountain forests of the Alps, assessments of their historical range of variability remain difficult. However, based on the increase of fire, bark beetle, and windthrow disturbances since the 20th century, which has coincided with a recovery of forest area and biomass, and based on analogies with similar ecosystems in Europe (Kulakowski et al., 2017), we can assume that the importance of the natural disturbances, which has been dominated by human activities for centuries, is increasing and will be an important driver of mountain forest dynamics in the Alps in future decades.

## Acknowledgments

We thank Laura Ramstein, Christian Rechsteiner and Carla Eschmann for research assistance. For providing data, we thank Davide Ascoli, Herbert Borchert, Beat Forster, Olivier Nägele, Boris Pezzatti, Marco Vanoni and Benjamin Zweifel. Dominik Kulakowski was supported by a fellowship from the Swiss Federal Institute for Forest, Snow and Landscape Research, WSL and R. Seidl acknowledges funding from an Austrian Science Fund FWF START grant (Y895-B25). We thank Thomas Thieke, Tom Nagel and two anonymous reviewers for helpful comments on an earlier version of the manuscript.

## Appendix A

See Table A1.

**Table A1**

Compilation of available data on occurrence, damage volume and size of disturbed forest patches in the Alps since the 19th century. The compilation is based on different and heterogenous sources and cannot be expected to be complete.

Agent	Region available	Data for	Affected forest	Other characteristics	Source
Wind	Switzerland <sup>d</sup>	1865–2014	0.16 mio m <sup>3</sup> /yr	26 winterstorms, 634 events	Usbeck (2015)
Wind	Swiss Alps	1990	4.5 mio m <sup>3</sup>	Max. 59.6 ha, median 0.5 ha	WSL Geodatabase
Wind	Bavarian Alps	1990	3.2 mio m <sup>3</sup>	ca. 3% of volume	König (1995)
Wind	Slovenian Alps	1918–2014	0.02 mio m <sup>3</sup> /yr		Bleiweis (1983)

Agent	Region available	Data for	Affected forest	Other characteristics	Source
Wind	Austrian Alps	1829	0.7 mio m <sup>3</sup>		Weigl (1997)
Wind	Switzerland <sup>a</sup>	1967	2.9 mio m <sup>3</sup>	0.9% of volume	Usbeck (2015)
Wind	Slovenian Alps	1984	0.45 mio m <sup>3</sup> /yr		Papler-Lampe (2008)
Wind	Swiss Alps	1999	5.6 mio m <sup>3</sup>	Max. 26 ha, 39% scattered <sup>c</sup>	WSL Geodatabase
Wind	Slovenian Alps	2008	0.3 mio m <sup>3</sup>	Over 9260 ha total area	SFS (2015)
Wind	Austrian Alps	2002–2010	1.9 mio m <sup>3</sup> /yr		Thom et al. (2013)
Wind and Snow	Austria <sup>a</sup>	1944–1989	1.6 mio m <sup>3</sup> /yr		Steyrer et al. (2011)
Wind and Snow	Aosta/Italian Alps	1990–2010	55 ha/yr	Max. 147.2 ha, median 0.1 ha	Vacchiano et al. (2016)
Snow	Switzerland <sup>a</sup>	1885	0.2 mio m <sup>3</sup>	Above 700 m asl	Coaz (1889)
Snow/Ice	Slovenian Alps	1918–2014	0.03 mio m <sup>3</sup> /yr		Zupancic (1969), SFS (2015)
Snow	Austrian Alps	1930–1931	0.4 mio m <sup>3</sup> /yr	Mainly 600–900 m asl	Weigl (1997)
Snow	Graubünden, CH	1990–2015	0.02 mio m <sup>3</sup> /yr	Mainly small scaled	Database Kt. Graubünden
Snow	Austrian Alps	2002–2010	0.3 mio m <sup>3</sup> /yr		Thom et al. (2013)
Snow/Ice	Slovenian Alps	2014	1.5 mio m <sup>3</sup>	Mainly broadleaved trees	SFS (2015)
Avalanches	Swiss Alps	1988	0.08 mio m <sup>3</sup>	1340 ha destroyed forest	Coaz (1889)
Avalanche	Swiss Alps	1891–2015	0.05 mio m <sup>3</sup> /yr	Max. 93 ha, median 1 ha	SLF-Avalanche database
Avalanches	Swiss Alps	1951	0.18 mio m <sup>3</sup>	2100 ha destroyed forest	SLF (1999)
Avalanches	Slovenian Alps	1951–1952	0.02 mio m <sup>3</sup>	450 ha destroyed forests	Cihal and Valinger (1959)
Avalanches	Swiss Alps	1999	0.16 mio m <sup>3</sup>	1400 ha destroyed forest	SLF (1999)
Bark beetles	Paneveggio/It	1920–1925	>0.1 mio m <sup>3</sup>	>30 years of prescribed cut	Motta et al. (2006a)
Bark beetles	Austria <sup>a</sup>	1944–1991	0.2 mio m <sup>3</sup> /yr		Steyrer et al. (2011)
Bark beetles	Switzerland <sup>a</sup>	1945–1950	0.25 mio m <sup>3</sup> /yr	On Spruce and Fir	Pfister (1988)
Bark beetles	Switzerland <sup>a</sup>	1988–2000	0.2 mio m <sup>3</sup> /yr		Meier et al. (2010)
Bark beetles	Switzerland <sup>a</sup>	2001–2007	1 mio m <sup>3</sup> /yr	Following blowdown in 1999	Meier et al. (2010)
Bark beetles	Austrian Alps	2002–2010	1.3 m <sup>3</sup> /yr		Thom et al. (2013)
Bark beetles	Aosta/Italian Alps	1984–2006 <sup>b</sup>	840 ha/yr	0.85 ha/yr	Vacchiano et al. (2016)
Fire	Northern Swiss Alps	1990–2014	2.5 ha/yr	8 fires/yr, median size 0.02 ha	Pezzatti et al. (2010)
Fire	Central Swiss Alps	1990–2014	33.6 ha/yr	22 fires/yr, median size 0.1 ha	Pezzatti et al. (2010)
Fire	Southern Swiss Alps	1990–2014	176.7 ha/yr	52.2 fires/yr, median size 0.1 ha	Pezzatti et al. (2010)
Fire	Aosta/ Italian Alps	1961–2010	147 ha/yr	Yearly 0.15% of total forest cover	Vacchiano et al. (2016)
Fire	Hautes Alpes/Fr	1976–2015	87.5 ha/yr	16 fires/yr, median size 1 ha	<a href="http://www.promethee.com">http://www.promethee.com</a>
Fire	Haute Provence/Fr <sup>a</sup>	1976–2015	374.7 ha/yr	39 fires/yr, median size 1 ha	<a href="http://www.promethee.com">http://www.promethee.com</a>
Fire	Alpes Maritimes/Fr <sup>a</sup>	1976–2015	1322.5 ha/yr	180 fires/yr, median size 0.3 ha	<a href="http://www.promethee.com">http://www.promethee.com</a>

<sup>a</sup>Area partly outside of Alps.

<sup>b</sup>Without years 1991–93.

<sup>c</sup>Scattered disturbance, 20–60% remaining crown cover of affected stands.

## References

- Abegg, M., Brändli, U-B., Cioldi, F., Fischer, C., Herold-Bonardi, A., Huber, M., Keller, M., Meile, R., Rösler, E., Speich, S., Traub, B., et al. Viertes Schweizerisches Landesforstinventar – Ergebnistabellen und Karten des LFI 2009-2013 (LFI4b). Birmensdorf: Eidg. Forschungsanstalt WSL; 2014. <<http://www.lfi.ch/resultate/>> [Published online 06.11.2014] [accessed 10.02.16]
- Ammer C. Impact of ungulates on structure and dynamics of natural regeneration of mixed mountain forests in the Bavarian Alps. *For Ecol Manage.* 1996; 88:43–53.
- Arpaci A, Malowerschnig B, Sass O, Vacik H. Using multi variate data mining techniques for estimating fire susceptibility of Tyrolean forests. *Appl Geogr.* 2015; 53:258–270.
- Ascoli D, Castagneri D, Valsecchi C, Conedera M, Bovio G. Post-fire restoration of beech stands in the Southern Alps by natural regeneration. *Ecol Eng.* 2013; 54:210–217.
- Arndt NA, Vacik HA, Koch VA, Arpaci AA, Gossow H. Modeling human-caused forest fire ignition for assessing forest fire danger in Austria. *iForest – Biogeosci Forest.* 2013; 6:315–325.
- AWLN. Iechtensteinisches Landeswaldinventar. Ergebnisse der dritten Erhebung 2010. Amt für Wald, Natur und Landschaft; Vaduz: 2012. p. 54
- Baldock, D., Beaufoy, G., Brouwer, F. Farming at the margins: abandonment or redeployment of agricultural land in Europe Case studies. Institute for European Environmental Policy (IEEP); The Hague: 1996. p. 202
- Bätzing, W. Die Alpen - Geschichte und Zukunft einer europäischen Kulturlandschaft. C.H. Beck; München: 2003.
- Bebi P, Kulakowski D, Rixen C. Snow avalanche disturbances in forest ecosystems – state of research and implications for management. *For Ecol Manage.* 2009; 257:1883–1892.
- Bebi P, Putallaz J-M, Fankhauser M, Schmid U, Schwitter R, Gerber W. Die Schutzfunktion in Windwurfflächen. *Schweizerische Zeitschrift für Forstwesen.* 2015; 166:168–176.
- BFW. Austrian Research Centre for Forests; 2016. Austrian forest inventories. <<http://bfw.ac.at/rz/wi.home>> [accessed 10. 02. 2016]
- Blarquez O, Carcaillet C, Bremond L, Maurier B, Radakovitch O. Trees in the subalpine belt since 11700 ca. BP: origin, expansion and alteration in the modern forest. *Holocene.* 2009; doi: 10.1177/0959683609348857
- Bleiweis, S. Pogostost in obseg škod zaradi ujm v slovenskih gozdovih. *Gozdarski vestnik*; 1983. p. 233-249.(in Slovene)
- Bork, H-R., Dalchow, C., Faust, B., Pierr, H-P., Toussaint, V., Werner, A. Future development of landscapes in marginal agrarian regions of central Europe: long-term effects of land use on the water balance. In: Tenhunen, JD.Lentz, R., Hantschel, R., editors. *Ecosystem Approaches to Landscape Management in Central Europe, Ecological Studies 147.* Berlin Heidelberg; New York: 2001. p. 571-581.
- Bottero A, Garbarino M, Long JN, Motta R. The interacting ecological effects of large-scale disturbances and salvage logging on montane spruce forest regeneration in the western European Alps. *For Ecol Manage.* 2013; 292:19–28.
- Buma B. Disturbance interactions: characterization, prediction, and the potential for cascading effects. *Ecosphere.* 2015; 6:1–15.
- Buma B, Wessman CA. Forest resilience, climate change, and opportunities for adaptation: a specific case of a general problem. *For Ecol Manage.* 2013; 306:216–225.
- Bunce RGH. Ecological implications of land abandonment in Britain: some comparison with Europe. *Options Méditerran.* 1991; 15:53–59.
- Büntgen U, Frank DC, Liebhold A, Johnson D, Carrer M, Urbinati C, Grabner M, Nicolussi K, Levanić T, Esper J. Three centuries of insect outbreaks across the European Alps. *New Phytol.* 2009; 182:929–941. [PubMed: 19383093]
- Cada V, Morrissey RC, Michalova Z, Bace R, Janda P, Svoboda M. Frequent severe natural disturbances and non-equilibrium landscape dynamics shaped the mountain spruce forest in central Europe. *For Ecol Manage.* 2016:363–178.

- Castebrunet H, Eckert N, Giraud G, Durant Y, Morin S. Projected changes of snow conditions and avalanche activity in a warming climate: the French Alps over the 2020–2050 and 2070–2100 periods. *Cryosphere*. 2014; 8:1673–1697.
- Cesti G, Conedera M, Spinedi F. Considerazioni sugli incendi boschivi causati dai fulmini. *Schweizerische Zeitschrift für Forstwesen*. 2005; 156:353–361.
- Chatré B, Lanzinger G, Macaluso M, Mayrhofer W, Morandini M, Onida M, Polajnar B. The Alps: People and pressures in the mountains, the facts at a glance. Permanent Secretariat of the Alpine Convention. 2010
- Chauchard S, Beilhe F, Denis N, Carcaillet C. An increase in the upper tree-limit of silver fir (*Abies alba* Mill) in the Alps since the mid-20th century: a land-use change phenomenon. *For Ecol Manage*. 2010; 259:1406–1415.
- Cihal, V., Valinger, E. Nekaj o sneznih plazovih. *Gozdarski vestnik*; 1959. p. 35-38.(in Slovene)
- Coaz, J. Der Schneeschaden vom 28./29. September 1885 in den Waldungen der Schweiz. *Stämpfli'sche Buchdruckerei*; Bern: 1887. p. 47(in German)
- Coaz, J. Der Lauinenschaden im schweizerischen Hochgebirge im Winter und Frühjahr 1887-88. *Stämpfli'sche Buchdruckerei*; Bern: 1889. p. 67
- Conedera M, Cesti G, Pezzatti GB, Zumbrunnen T, Spinedi F, Viegas DX. Lightning-induced fires in the Alpine region: an increasing problem. V. Int. Conference on Forest Fire Research. 2006
- Conedera M, Peter L, Marxer P, Forster F, Rickenmann D, Re L. Consequences of forest fires on the hydrogeological response of mountain catchments: a case study of the Riale Buffaga, Ticino, Switzerland. *Earth Surf Proc Land*. 2003; 28:117–129.
- Conedera M, Tonini M, Oleggini L, Orozco CV, Leuenberger M, Pezzatti GB. Geospatial approach for defining the Wild land-Urban Interface in the Alpine environment. *Comput Environ Urban Syst*. 2015; 52:10–20.
- Conedera M, Colombaroli D, Tinner W, Krebs P, Whitlock C. Insights about past forest dynamics as a tool for present and future forest management in Switzerland. *Forest Ecol Manage*. 2017; 388:100–112.
- Dale VH, Joyce LA, McNulty S, Neilson RP, Ayres MP, Flannigan MD, et al. Climate change and forest disturbances. *Bioscience*. 2001; 51:723–734.
- Dambrine E, Dupouey J-L, Laüt L, Thinon M, Beaufiles T, Richard R. Present forest biodiversity patterns in France related to former roman agriculture. *Ecology*. 2006; 88:1430–1439.
- Deankovic T. Snegolomi v Julijskih Alpah in njihovi vzroki. *Gozdarski vestnik*. 1969; 27(9):223–236. (in Slovene).
- Didion M, Kupferschmid AD, Bugmann H. Long-term effects of ungulate browsing on forest composition and structure. *For Ecol Manage*. 2009; 258:44–55.
- Dobbertin M. Influence of stand structure and site factors on wind damage comparing the storms Vivian and Lothar. *Forest, Snow Landscape Res*. 2002; 77:187–205.
- Flatley W, Lafon CW, Grissino-Mayer HD, Forest LB. Fire history, related to climate and land use in three southern Appalachian landscapes in the eastern United States. *Ecol Appl*. 2013; 23:1250–1266. [PubMed: 24147399]
- Flinn KM, Vellend M. Recovery of forest plot communities in post-agricultural landscapes. *Front Ecol*. 2005; 3:243–250.
- Föhn, P. Avalanche frequency and risk estimation in forest sites. *Proc. IUFRO Seminar Mountain Forests and Avalanches*; Sept. 1978; Davos. 1978. p. 242-254.
- Foster DR. Land-use history (1730–1990) and vegetation dynamics in Central New England, USA. *J Ecol*. 1992; 80:753–771.
- Foster DR, Swanson F, Aber J, Burke I, Brokaw N, Tilman D, Knapp A. The importance of land-use legacies to ecology and conservation. *Bioscience*. 2003; 53:78–88.
- Garbarino M, Lingua E, Subira MM, Motta R. The larch wood pasture: structure and dynamics of a cultural landscape. *Eur J Forest Res*. 2011; 130:491–502.
- Gardiner, B., Blennow, K., Carnus, JM., Fleischer, P., Ingemarson, F., Landmann, G., Lindner, M., Marzano, M., Nicoll, B., Orazio, C., Peyron, JL., et al. Destructive storms in European forests: past


- and forthcoming impacts. Destructive Storms in European Forests: Past and Forthcoming Impacts. Gardiner, B., et al., editors. European Forest Institute; Joensuu: 2010. p. 138
- Gehrig-Fasel J, Gusian A, Zimmermann NE. Tree line shifts in the Swiss Alps: climate change or land abandonment. *J Veg Sci.* 2007; 18:571–582.
- Gellrich M, Baur P, Koch B, Zimmermann N. Agricultural land abandonment and natural forest regrowth in the Swiss mountains: a spatially explicit economic analysis. *Agric Ecosyst Environ.* 2007; 118:93–108.
- Ginzler C, Brändli UB, Hägeli M. Waldflächenentwicklung der letzten 120 Jahre in der Schweiz. *Schweizerische Zeitschrift für Forstwesen.* 2011; 162:337–334.
- Hauser A. Wald und Forst in der Gründungszeit der Eidgenossenschaft. *Schweizerische Zeitschrift für Forstwesen.* 1964; 115:637–658.
- Hlasny T, Kristek S, Holusa J, Trombik J, Urbancova N. Snow disturbances in secondary Norway spruce forests in Central Europe: regression modeling and its implications for forest management. *For Ecol Manage.* 2011a; 262:2151–2161.
- Hlasny T, Zajickova L, Turcani M, Holusa J, Sitkova Z. Geographical variability of spruce bark beetle development under climate change in the Czech Republic. *J For Sci.* 2011b; 57:242–249.
- Janda P, Ba e R, Trotsiuk V, Mikoláš M, Nagel T, Seidel R, Mrhalová H, Morrissey RC, Matěj L, Lábusová J, Kuchel S, et al. The historical disturbance regime of mountain Norway spruce forests in the Western Carpathians and its influence on current forest structure and composition. *For Ecol Manage.* 2017; 388:67–8.
- Kaplan JO, Krumhardt KM, Zimmermann N. The prehistoric and preindustrial deforestation of Europe. *Quatern Sci Rev.* 2009; 28:3016–3034.
- Keller, M. Schweizerisches Landesforstinventar. Anleitung für die Felddaufnahmen der Erhebung 2004–2007. WSL; Birmensdorf: 2005.
- Klopčič M, Poljanec A, Gartner A, Boncina A. Factors related to natural disturbances in mountain Norway spruce (*Picea abies*) forests in the Julian Alps. *Ecoscience.* 2009; 16:48–57.
- Klopčič M, Boncina A. Stand dynamics of silver fir (*Abies alba* Mill.) – European beech (*Fagus sylvatica* L.) forests during the past century: a decline of silver fir? *Forestry.* 2011; 84:259–271.
- König, A. Sturmgefährdung von Beständen im Altersklassenwald. J.D. Sauerländer's Verlag; Frankfurt am Main: 1995.
- Körner W, Dupouey E, Dambrine E, Benoit M. Influence of past land use on the vegetation and soils of present day forest in the Vosges mountains. *J Ecol.* 1997; 85:351–358.
- Kozak J. Forest cover change in the Western Carpathians in the past 180 years. *Mountain Res Develop.* 2003; 23:369–375.
- Kral F. Nacheiszeitliche Waldentwicklungstypen in den Alpen. *Verh Zool-Bot Ges Österreich.* 1995; 132:1–12.
- Kraus, H., Ebel, U. Risiko Wetter: Die Entstehung von Stürmen und anderen atmosphärischen Gefahren. Springer; Berlin: 2003.
- Krumm F, Kulakowski D, Risch AC, Spiecker H, Brändli U-B, Bebi P. Stem exclusion and mortality in unmanaged subalpine forests of the Swiss Alps. *Eur J Forest Res.* 2012; 131:1571–1583.
- Kulakowski D, Veblen TT. Influences of fire history and topography on the pattern of a severe wind blowdown in a Colorado subalpine forest. *J Ecol.* 2002; 90:806–819.
- Kulakowski D, Veblen TT, Bebi P. Effects of fire and spruce beetle outbreak legacies on the disturbance regime of a subalpine forest in Colorado. *J Biogeogr.* 2003; 30:1445–1456.
- Kulakowski D, Bebi P. Range of variability of unmanaged subalpine forests. *Forum für Wissen.* 2004; 2004:47–54.
- Kulakowski D, Bebi P, Rixen C. The interacting effects of land use change, climate change and suppression of natural disturbances on landscape forest structure in the Swiss Alps. *Oikos.* 2011; 120:216–225.
- Kulakowski D, Jarvis D, Veblen TT, Smith J. Stand-replacing fires reduce susceptibility to mountain pine beetle outbreaks in Colorado. *J Biogeogr.* 2012; 39:2052–2060.

- Kulakowski D, Matthews C, Jarvis D, Veblen TT. Compounded disturbances in subalpine forests in western Colorado favor future dominance by quaking aspen (*Populus tremuloides*). *J Veg Sci*. 2013; 24:168–176.
- Kulakowski D, Seidl R, Holeksa J, Kuuluvainen T, Nagel T, Panayotov M, Svoboda M, Thorn S, Vacchiano G, Whitlock C, Wohlgemuth T, et al. A walk on the wild side: disturbance dynamics and the conservation and management of European mountain forest ecosystems. *Forest Ecol Manage*. 2017; 388:120–131.
- Küster, H. Geschichte der Landschaft in Mitteleuropa: Von der Eiszeit bis zur Gegenwart. C.H. Beck; München: 2010. p. 448
- Küttel M. The subalpine protection forest in the Urseren valley – an inelastic ecosystem. *Bot Helv*. 1990; 100:183–197.
- Landolt, E. Bericht an den hohen schweizerischen Bundesrat über die Untersuchung der schweiz. Hochgebirgswaldungen, vorgenommen in den Jahren 1858 1859 und 1860. Bern: 1862.
- Latenser, M., Pfister, C. Avalanches in Switzerland 1500–1990. Rapid Mass Movement as a Source of Climatic Evidence for the Holocene. Frenzel, B., et al., editors. 1997. p. 241–266.
- Liu Z, Wimberly MC, Lamsal A, Sohl TL, Hawbaker TJ. Climate change and wildfire risk in an expanding wildland–urban interface: a case study from the Colorado Front Range Corridor. *Landscape Ecol*. 2015; 30:1943–1957.
- Mather AS, Fairbairn J. From floods to reforestation: the forest transition in Switzerland. *Environ History*. 2000; 6:399–421.
- Mather AS, Fairbairn J, Needle CL. The course and drivers of the forest transition: the case of France. *J Rural Stud*. 1999; 15:65–90.
- Mathieu, J., Backhaus, N., Hürlimann, K., Bürgi, M. Geschichte der Landschaft in der Schweiz. Orell Füssli; Zürich: 2016. p. 384
- McEvedy, C., Jones, R. Atlas of World Population History. Penguin Books Ltd; Harmondsworth: 1978.
- Meier, F., Forster, B., Engesser, R. Waldschutz aktuell 01/2001 bis 01/2009. 2010. <[http://www.wsl.ch/fe/walddynamik/waldschutz/wsinfo/wsaktuell\\_DE](http://www.wsl.ch/fe/walddynamik/waldschutz/wsinfo/wsaktuell_DE)>
- Moser B, Temperli C, Schneiter G, Wohlgemuth T. Potential shift in tree species composition after interaction of fire and drought in the Central Alps. *Eur J Forest Res*. 2010; 129:625–633.
- Motta R, Berretti R, Lingua E, Piussi P. Coarse woody debris, forest structure and regeneration in the Valbona Forest Reserve, Paneveggio, Italian Alps. *For Ecol Manage*. 2006a; 235:155–163.
- Motta R, Morales M, Nola P. Human land-use, forest dynamics and tree growth at the treeline in the Western Italian Alps. *Ann For Sci*. 2006b; 63:739–747.
- Müller MM, Vacik H, Diendorfer G, Arpacı A, Formayer H, Gossow H. Analysis of lightning-induced forest fires in Austria. *Theoret Appl Climatol*. 2012; 111:183–193.
- Netherer S, Matthews B, Katzensteiner K, Blackwell E, Henschke P, Hietz P, Pennerstorfer J, Rosner S, Kikuta S, Schume H, Schopf A. Do water-limiting conditions predispose Norway spruce to bark beetle attack? *New Phytol*. 2015; 205:1128–1141. [PubMed: 25417785]
- Nykänen ML, Peltola H, Quine C, Kellomäki S, Broadgate M. Factors affecting snow damage of trees with particular reference to European conditions. *Silva Fennica*. 1997; 31:193–213.
- Paine RT, Tegner MJ, Johnson EA. Compounded perturbations yield ecological surprises. *Ecosystems*. 1998; 16:535–545.
- Panayotov M, Gogushev G, Tsavkov E, Vassileva P, Tsvetanov N, Kulakowski D, Bebi P. Abiotic disturbances in Bulgarian mountain coniferous forests – an overview. *For Ecol Manage*. 2017; 388:13–28.
- Papler-Lampe V. Snegolom, ki je januarja 2007 prizadel blejske gozdove. *Gozdarski vestnik*. 2008; 66(5–6):309–19. (In Slovene).
- Patzelt G, Bortenschlager S. Die postglazialen Gletscher- und Klimaschwankungen in der Venedigergruppe (Hohe Tauern, Ostalpen). *Zeitschrift für Geomorphologie N.F Suppl*. Bd. 1973; 16:25–72.
- Paulsen J, Körner C. GIS-analysis of tree-line elevation in the Swiss Alps suggests no exposure effect. *J Veg Sci*. 2009; 12:817–824.


- Petek, F. Spremembe rabe tal v slovenskem alpskem svetu Geografija Slovenije 11. Ljubljana: Založba ZRC; 2005. p. 216(in Slovene)
- Pezzatti GB, Reinhard M, Conedera M. Swissfire: die neue schweizerische Wald-branddatenbank. Schweiz Z Forstwes. 2010; 161:465–469.
- Pfister, C. Bundesamt für Forstwesen und Landschaftsschutz; Bern: 1988. Witterungsextreme und Waldschäden in der Schweiz: eine historisch-kritische Untersuchung von Schadenmeldungen aus schweizerischen Wäldern in ihrer Beziehung zur Klimabelastung, insbesondere durch sommerliche Dürreperioden.
- Plue J, Hermy M, Verheyen K, Thuillier P, Saguez R, Decoq G. Persistent changes in forest vegetation and seed bank 1600 years after human occupation. Landscape Ecol. 2008; 23:673–688.
- Price M. Legislation and policy for forests of the Swiss Alps. Land Use Policy. 1988; 5:314–328.
- Puerta-Piñero C, Espelta JM, Sánchez-Humanes B, Rodrigues A, Coll L, Brotons L. History matters: previous land use changes determine post-fire vegetation recovery in forested Mediterranean landscapes. For Ecol Manage. 2012; 279:121–127.
- R Core Team. R Foundation for Statistical Computing; Vienna, Austria: 2013. R: A Language and Environment for Statistical Computing. <<http://www.R-project.org>>
- Rottmann, M. J.D. Sauerländers Verlag; Frankfurt: 1985. Schneebruchschäden in Nadelholzbeständen.
- Rozman A, Diaci J, Krese A, Fidej G, Rozenbergar D. Forest regeneration dynamics following bark beetle outbreak in Norway spruce stands: Influence of meso-relief, forest edge distance and deer browsing. Forest Ecol Manage. 2015; 353:196–207.
- Seidl R. The shape of ecosystem management to come: anticipating risks and fostering resilience. Bioscience. 2014; 64:1159–1169. [PubMed: 25729079]
- Seidl R, Schelhaas MJ, Lindner M, Lexer MJ. Modelling bark beetle disturbances in a large scale forest scenario model to assess climate change impacts and evaluate adaptive management strategies. Reg Environ Change. 2009; 9:101–119.
- Seidl R, Rammer W. Climate change amplifies the interactions between wind and bark beetle disturbance in forest landscapes. Landscape Ecol. in press.
- Seidl R, Schelhaas MJ, Lexer M. Unraveling the drivers of intensifying forest disturbance regimes in Europe. Global Change Biol. 2011a; 17:2842–2852.
- Seidl R, Rammer W, Lexer MJ. Climate change vulnerability of sustainable forest management in the Eastern Alps. Clim Change. 2011b; 106:225–254.
- Seidl R, Schelhaas M-J, Rammer W, Verkerk PJ. Increasing forest disturbances in Europe and their impact on carbon storage. Nat Clim Change. 2014; 4:806–810.
- Seidl R, Müller J, Hothorn T, Bässler C, Heurich M, Kautz M. Small beetle, large-scale drivers: how regional and landscape factors affect outbreaks of the European spruce bark beetle. J Appl Ecol. 2016; 53:530–540.
- Schelhaas M-J, Nabuurs GJ, Schuck A. Natural disturbances in the European forests in the 19th and 20th centuries. Glob Change Biol. 2003; 9:1620–1633.
- Schmucki E, Marty C, Fierz C, Weingartner R, Lehning M. Impact of climate change in Switzerland on socioeconomic snow indices. Theor Appl Climatol. 2015; doi: 10.1007/s00704-015-1676-7
- SFS. Forest Inventory Database and Regional Forest Management Plans (2011–2020). Slovenia Forest Service; Ljubljana, Slovenia: 2012.
- SFS. Database and Reports. Slovenia Forest Service; Ljubljana, Slovenia: 2015.
- SLF. Der Lawinenwinter. Ereignisanalyse. Davos, WSL-Institut für Schnee und Lawinenforschung; 1999. p. 588
- Spohn M, Novák T, Incze J, Giani L. Dynamics of soil carbon, nitrogen, and phosphorus in calcareous soils after land-use abandonment—a chronosequence study. Plant Soil. 2016; 401:185–196.
- Schuler, A. Forest area and forest utilisation in the Swiss pre-alpine region. In: Salbitano, F., editor. Human Influences on Forest Ecosystems Development in Europe. Pitagora Editrice; Bologna: 1988. p. 121–127.
- Schwoerer C, Colombaroli D, Kaltenrieder P, Rey F, Tinner W. Early human impact (5000–3000 BC) affects mountain forest dynamics in the Alps. J Ecol. 2015; 103:281–295.

- Sonnier J. Analyse du rôle de protection des forêts domaniales de montagne. *Revue Forestière Française*. 1991; 43:131–145.
- Stadelmann G, Bugmann H, Wermelinger B, Bigler C. Spatial interactions between storm damage and subsequent infestations by the European spruce bark beetle. *For Ecol Manage*. 2014; 318:167–174.
- Stahli M, Finsinger W, Tinner W, Allgoewer B. Wildfire history and fire ecology of the Swiss National Park (Central Alps): new evidence from charcoal, pollen and plant macrofossils. *Holocene*. 2006; 16:805–817.
- Steyrer G, Krenmayer W, Schaffer H. Dokumentation der Waldschädigungsfaktoren (DWF). *Forstschutz Aktuell*. 2011; 52:21–24.
- Temperli C, Bugmann H, Elkin C. Cross-scale interactions among bark beetles, climate change, and wind disturbances: a landscape modeling approach. *Ecol Monogr*. 2013; 83:383–402.
- Tinner W, Hubschmid P, Wehrli M, Ammann B, Conedera M. Long-term forest fire ecology and dynamics in southern Switzerland. *J Ecol*. 1999; 87:273–289.
- Tinner W, Kaltenrieder P. Rapid responses of high-mountain vegetation to early Holocene environmental changes in the Swiss Alps. *J Ecol*. 2005; 93:936–947.
- Tinner W, Conedera M, Ammann B, Lotter AF. Fire ecology north and south of the Alps since the last ice age. *Holocene*. 2005; 15:1214–1226.
- Thom D, Seidl R, Steyrer G, Krehan H, Formayer H. Slow and fast drivers of the natural disturbance regime in Central European forest ecosystems. *For Ecol Manage*. 2013; 307:293–302.
- Thom D, Rammer W, Seidl R. Disturbances catalyze the adaptation of forest ecosystems to changing climate conditions. *Glob Change Biol*. 2016; in press. doi: 10.1111/gcb.13506
- Usbeck T, Wohlgemuth T, Pfister C, Volz R, Beniston M, Dobbertin M. Wind speed measurements and forest damage in Canton Zurich (Central Europe) from 1891 to winter 2007. *Int J Climatol*. 2010a; 30:347–358.
- Usbeck T, Wohlgenuth T, Dobbertin M, Pfister C, Bürgi A, Rebetez M. Increasing storm damage to forests in Switzerland from 1858 to 2007. *Agric For Meteorol*. 2010b; 150:47–55.
- Usbeck T, Waldner P, Dobbertin M, Ginzler C, Hoffmann C, Sutter F, Steinmeier C, Volz R, Schneiter G, Rebetez M. Relating remotely sensed forest damage data to wind data: storms Lothar (1999) and Vivian (1990) in Switzerland. *Theoret Appl Climatol*. 2012; 108:451–462.
- Usbeck T. Wintersturmschäden im Schweizer Wald von 1865 bis 2014. *Schweizerische Zeitschrift für Forstwesen*. 2015; 166:184–190.
- Veblen TT, Hadley KS, Nel EM, Kitzberger T, Reid M, Villalba R. Disturbance regime and disturbance interactions in a Rocky Mountain Subalpine Forest. *J Ecol*. 1994; 82:125–135.
- Vacchiano, G., Dobbertin, M., Egli, S., Giordano, L., Gonthier, P., Mazzoglio, P.J., Motta, R., Nola, P., Nicolotti, G., Patetta, A., Polomski, J., et al. Il deperimento del pino silvestre nelle Alpi occidentali: natura ed indirizzi di gestione. *Compagnia delle Foreste*; Arezzo: 2008.
- Vacchiano G, Maggioni M, Perseghin G, Motta R. Effect of avalanche frequency on forest ecosystem services in a spruce-fir mountain. *Cold Reg Sci Technol*. 2015; 115:9–21.
- Vacchiano G, Berretti R, Mondino EB, Meloni F, Motta R. Assessing the effect of disturbances on the functionality of direct protection forests. *Mountain Res Develop*. 2016; 36:41–55.
- Vacik H, Arndt N, Arpacı A, Koch V, Müller M, Gossov H. Characterisation of forest fires in Austria. *Austrian J For Sci*. 2011; 2011:1–32.
- Valese E, Conedera M, Held AC, Ascoli D. Fire, humans and landscapes in the European Alpine region during the Holocene. *Anthropocene*. 2014; 2014:63–74.
- van der Knaap WO, van Leeuwen JFN, Finsinger W, Gobet E, Pini R, Schweizer A, Valsecchi V, Ammann B. Migration and population expansion of *Abies*, *Fagus*, *Picea*, and *Quercus* since 15000 years in and across the Alps, based on pollen-percentage threshold values. *Quatern Sci Rev*. 2005; 24:645–680.
- Wastl C, Schunk C, Lupke M, Cocca G, Conedera M, Valese E, Menzel A. Large-scale weather types, forest fire danger, and wildfire occurrence in the Alps. *Agric For Meteorol*. 2013; 168:15–25.
- Wear DN, Bolstad P. Land-use changes in southern Appalachian landscapes: spatial analysis and forecast evaluation. *Ecosystems*. 1998; 1:575–594.


- Weigl, N. Österreichs Forstwirtschaft in der Zwischenkriegszeit 1918–1938. Österreichischer Kunst- und Kulturverlag; Wien: 1997. p. 219
- Westerling AL, Hidalgo HG, Cayan DR, Westman TW. Warming and earlier spring increase western U.S. forest wildfire activity. *Science*. 2006; 313:940–943. [PubMed: 16825536]
- Winckler, K. Die Alpen im Frühmittelalter. Die Geschichte eines Raumes in den Jahren 500–800. Böhlau Verlag; Wien: 2012.
- Wohlgemuth T, Bürgi M, Scheidegger C, Schütz M. Dominance reduction of species through disturbance – a proposed management principle for Central European forests. *Forest Ecol Manage*. 2002; 166:1–15.
- Zumbrunnen T, Bugmann H, Conedera M, Bürgi M. Linking forest fire regimes and climate – a historical analysis in a dry inner Alpine valley. *Ecosystems*. 2009; 12:73–86.
- Zupancic M. Vetrolomi in snegolomi v Sloveniji v povojni dobi. *Gozdarski vestnik*. 1969:193–210. in Slovene.




**Fig. 1.** Examples of photographic time series visualizing forest cover changes in the Alps since the 19th century: (A and B): Ceüse, southern French Alps, at the end of the 19th century (A) at the beginning of the 21st century after so-called RTM (mountain terrains rehabilitation) works (B); (C and D): Davos (Central Swiss Alps) in 1900 (C) and 2010 (D); (E and F): Vermiglio (Trentino, Italian Alps) in 1915 (E) and 2000 (F). -Source: Trento Autonomous Province archive.




**Fig. 2.** Average annual burnt area in ha per year between (1985 and 2011) for different regions of the European Alps. Original data source: European Forest Fire Information System (EFFIS), Valese et al. (2014) and Vacik et al. (2011).




**Fig. 3.** Forest cover changes in different regions of the Alps since the availability of data. Black bars represent for each region the area of forest cover during the last available survey relative to the area of forest cover during the first available forest cover data (grey bars, standardized to same bar size for each region). Note that the time for the first available survey varies considerably between 1827 (Slovenian Alps; Petek, 2005) and 1981 (Piedmont).

**Fig. 4.**

Increase of forest cover percentage since 1880 as a function of distances from potential treeline (calculated based on moving window approach of the highest trees in a 10 km neighborhood). Fig. 4a shows forest cover percentage in 1880, 1940 and 2000 based on digitized historic maps. Fig. 4b shows relative increases between 1880 and 1940 (black bars) and between 1940 and 2000 (grey bars) as a ratio of original forest cover percentage.



**Fig. 5.** NFI-plots windthrown by the storms Vivian (1990) and Lothar (1999), established prior to 1880 (blue dots,  $n = 110$ ) and after 1880 (orange dots,  $n = 24$ ) in Switzerland. Barplots with corresponding colours represent the frequency of observed windthrow areas in pre- and post-1880 forests relative to the expected frequencies according to the area of each forest type.



**Fig. 6.** Representation of forest fire in NFI-plots which overlap with pre-1880 forests (blue dots,  $n = 236$ ) and post 1880-forests (orange dots, 193 Fires) in the Canton Graubünden (eastern Switzerland). Barplots with corresponding colours represent the frequency of observed fire areas in pre- and post-1880 forests relative to the expected frequencies according to the area of each forest type.

**Table 1**

Change of wood volume (live and dead) and species composition during the last decades according to forest inventory data from Austria (BFW, 2016; 1992/96–2007/09), Switzerland (Abegg et al., 2014; 1993/95–2009/13), Slovenia (SFS, 2012; 1970–2010), France (1973–2009/13, uploaded from [www.inventaire-forestier-ign.fr](http://www.inventaire-forestier-ign.fr) on the 22th of April 2016) and Liechtenstein (AWLN, 2012, 1986–2010).

Region	Live volume		Dead volume		Change species share per decade (%)		
	[m <sup>3</sup> /ha] <sup>a</sup>	Change per decade (%)	[m <sup>3</sup> /ha] <sup>a</sup>	Change per decade (%)	Spruce	Fir	Beech
Vorarlberg (A)	411	+6.1	11.1	+50.7	+1.9	−2.6	−0.3
Tyrol (A)	328	+9.3	10.8	+60.7	−0.6	+0.2	+0.1
Salzburg (A)	346	+7.8	10.6	+58.7	−1.1	0.0	0.0
Carinthia (A)	352	+13.8	7.7	+49.8	−0.1	−0.2	+0.4
Styria (A)	352	+9.3	8.6	+47.5	−0.1	+0.1	+0.7
Liechtenstein	409	+2.4	30.0	NA	−5.8	−3.8	+2.5
Northern Swiss Alps	456	+0.4	36.5	105.8	−2.1	+0.6	+0.5
Central Swiss Alps	347	+7.7	32.3	+39.3	−1.1	+0.7	+0.3
Southern Swiss Alps	278	+17.1	22.5	+59.1	−0.9	0.0	+0.9
Northern French Alps	274	+12.0	NA	NA	+5.1	+11.0	+20.0
West. Slovenian Alps	313	+9.9	13.9	NA	−0.8	−1.0	+1.3
East. Slovenian Alps	352	+16.9	36.1	NA	+1.5	−2.6	+0.6

<sup>a</sup> Assessed at the last available inventory.

**Table 2**

Comparison of pre-1880 and post-1880-forests according to different attributes deduced from forest inventory data (NFI) for the Swiss Alps. Data pertain to observations in the observation period NFI4 4 (2009–2013).

Attributes	Pre-1880 forests Average value (Stdv.)	Post 1880 forests Average value (Stdv.)	p-values Wilcoxon rank-sum test
Total current wood volume	420.9 m <sup>3</sup> /ha (270)	343.8 m <sup>3</sup> /ha (274)	<0.001
Dead wood volume	31.3 m <sup>3</sup> /ha (57)	27.3 m <sup>3</sup> /ha (56.2)	0.02
Attributes	Pre-1880 forests Number of counts	Post 1880 forests Number of counts	p-values Chi-squared
Total number of plots	1114	856	
Young growth (<12 cm DBH)	66 (6%)	54 (6%)	n.s.
Pole wood (12–30 cm DBH)	156 (14%)	180 (21%)	<0.001
Timber wood 1 (30–50 cm DBH)	306 (27%)	202 (21%)	0.05
Timber wood 2 (>50 cm DBH)	275 (25%)	125 (15%)	<0.001
Mixed diameter class	311 (28%)	295 (34%)	0.002
Crown closure: closed	284 (25%)	202 (24%)	n.s.
Crown closure: normal-loose	331 (30%)	217 (25%)	0.04
Crown closure: scattered-open	295 (26%)	244 (29%)	n.s.
Crown closure: spatially aggregated	120 (11%)	159 (19%)	<0.001
Crown closure: multilayered dense	84 (8%)	34 (4%)	0.001
Dominant tree species <i>Picea abies</i>	540 (48%)	377 (44%)	0.02
Dominant tree species <i>Abies alba</i>	147 (13%)	24 (3%)	<0.001
Dominant tree species <i>Larix decidua</i>	96 (8%)	146 (15%)	<0.001
Other coniferous dominant species	63 (6%)	43 (3%)	n.s.
Dominant tree species <i>Fagus sylvatica</i>	147 (10%)	82 (13%)	0.02
Other broadleaved dominant species	124 (11%)	195 (22.8)	<0.001
Signs of grazing	78 (7%)	163 (19%)	<0.001
Last management intervention 20 years	616 (55%)	351 (41%)	<0.001
Last management intervention >50 years	316 (28%)	358 (43%)	<0.001

Note: n.s.: p-values >0.05.